



ERC Catalysts, Inc.
 215 Main Street, Miltona, MN
 Phone: 1.218.943.7904

HEMICELLULASE (*Aspergillus niger*)

SPECIFICATION SHEET

I.U.B. 3.2.1.78

DESCRIPTION

Hemicellulase is endo-enzyme and is derived from the fermentation of a selected Non-GMO str *Aspergillus niger*.

PHYSICAL PROPERTIES

An off white to light tan colored, free-flowing powder, soluble in water, free of offensive odor ;

ENZYMATIC PROPERTIES

The pH optimum is 4.0 with a stability range of 3.0 to 6.0. The optimum temperature is 55°C range of 50°C to 60°C.

ACTIVITY

The activity is characterized by its ability to hydrolyze galactomannoglucans, mannans and glu. The acceptance criteria for all enzyme assays is: NLT 85.0% and NMT 115.0% of the declared u enzyme activity.*

COUNTRY OF ORIGIN Japan

STORAGE/ SHELF LIFE/ STANDARD PACK SIZE

Product is stable for two years (24 months) if stored at or below 10°C in sealed poly bags in bc away from sunlight and high humidity. Product is packed in 25 kilo fiber drums or double-wall boxes.

HANDLING PRECAUTIONS

Avoid the formation of aerosol and dust of the product. Repeated inhalation of enzyme aerosc cause allergic type reactions in sensitized individuals. For detailed information please refer to

Description	Specification	Method
Activity:	NLT 400,000 HCU/GM	FCC
Identity:	Hemicellulase Fungal	FTIR
Moisture:	NMT 10%	Ohaus MB-45
Metals:		
Lead	NMT 5 ppm	SW-846 6020
Microbiological Data:		
TPC	<10,000 CFU/g	Soleris / AOAC 990.12
E. coli	Negative/10g	Soleris / AOAC 991.14
Enteroc	<100 CFU/g	AOAC 2003.01
Salmonella**	Negative/25g	BAM Ch. 5/AOAC 2011.03
Yeast	<1,000 CFU/g	Soleris / AOAC 997.02
Mold	<1,000 CFU/g	Soleris / AOAC 997.02
Coliforms	<100 CFU/g	Soleris / AOAC 991.14

*FCC 9 Page 414

**If Enteroc test results exceed 100 CFU/g then Salmonella testing is completed.