



ERC Catalysts, Inc.  
215 Main Street, Miltona, MN 56354 U.S.A  
Phone: 218.943.7904

## TRYPSIN (Porcine)

### SPECIFICATION SHEET

I.U.B. 3.4.21.4

### DESCRIPTION

Trypsin is derived from porcine pancreas. It is a proteolytic enzyme that is active in the alkaline pH range.

### PHYSICAL PROPERTIES

A white to off white colored, free-flowing powder, soluble in water, with characteristic odor and taste.

### ENZYMATIC PROPERTIES

The pH optimum range is 7.4 to 8.2. The optimum temperature range is 38°C to 40°C.

### ACTIVITY

The acceptance criteria for all enzyme assays is: NLT 85.0% and NMT 115.0% of the declared units of activity.\*

**COUNTRY OF ORIGIN** China

### STORAGE/SHELF LIFE/STANDARD PACK SIZE

Product is stable for two years (24 months) if stored at or below 10°C in sealed poly bags in boxes or drums away from sunlight and high humidity. Product is packed in 25 kilo fiber drums or double-wall bags.

### HANDLING PRECAUTIONS

Avoid the formation of aerosol and dust of the product. Repeated inhalation of enzyme aerosol or dust may cause allergic type reactions in sensitized individuals. For detailed information please refer to the SDS.

Description	Specification	Method
Activity:	NLT 2,400 USP/MG	USP
Identity:	Trypsin	FTIR
Moisture:	NMT 10%	Ohaus MB-45
Metals:		
Lead	NMT 5 ppm	SW-846 6020
Microbiological Data:		
TPC	<10,000 CFU/g	Soleris / AOAC 990.12
E.coli	Negative/10g	Soleris / AOAC 991.14
Enterococcus	<100 CFU/g	AOAC 2003.01
Salmonella**	Negative/25g	BAM Ch. 5 / AOAC 2011.03
Yeast	<1,000 CFU/g	Soleris / AOAC 997.02
Mold	<1,000 CFU/g	Soleris / AOAC 997.02
Coliforms	<100 CFU/g	Soleris / AOAC 991.14

\*FCC 9 Page 414

\*\*If Enterococcus test results exceed 100 CFU/g then Salmonella testing is completed.

ange.

te.

enzyme

all boxes.

1st may  
1955.